

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 962 452 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
17.09.2003 Bulletin 2003/38

(51) Int Cl.7: **C07D 207/09, C07D 207/12,
A61K 7/13**

(21) Application number: **99201486.0**

(22) Date of filing: **12.05.1999**

(54) **Oxidative hair dye compositions and methods containing
1-(4-aminophenyl)-2-pyrrolidinemethanols**

1-(4-Aminophenyl)-2-pyrrolidinemethanole enthaltende oxidative Haarfärbemittel und Verfahren

Composition de teinture d'oxydation et procédés comprenant de dérivés 2-pyrrolidinéméthanol
1-(4-aminophényle)

(84) Designated Contracting States:
DE FR GB IT SE

(30) Priority: **13.05.1998 US 78264**

(43) Date of publication of application:
08.12.1999 Bulletin 1999/49

(73) Proprietor: **P&G-Clairol, Inc.**
Stamford, CT 06922 (US)

(72) Inventors:
• **Lim, Mu-I11**
Trumbull, Connecticut 06611 (US)
• **Popp, Margaret**
Branford, Connecticut 06405 (US)
• **Pan, Yuh-Guo**
Stamford, Connecticut 06905 (US)

(74) Representative:
Adams, Harvey Vaughan John et al
Mathys & Squire,
100 Gray's Inn Road
London WC1X 8AL (GB)

(56) References cited:
EP-A- 0 634 164 **US-A- 5 278 034**

- **PINIELLA J F ET AL: "Synthesis, structure and second-order nonlinear optical properties of highly functionalized 6-aminopentafulvenes" J. MATER. CHEM. (JMACEP,09599428);1998; VOL.8 (3); PP.619-627, XP002116645 Univ. Autonoma de Barcelona;Barcelona; 08193; Spain (ES)**
- **PATENT ABSTRACTS OF JAPAN vol. 014, no. 430 (P-1106), 14 September 1990 (1990-09-14) & JP 02 167529 A (AJINOMOTO CO INC), 27 June 1990 (1990-06-27)**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 962 452 B1

Description

FIELD OF THE INVENTION

[0001] The invention relates generally to methods and compositions for preparing stable oxidative hair dyes that result in long-lasting and true colors which are different from colors produced by similar primary intermediates. The present invention more particularly relates to oxidative hair dye compositions and methods comprising 1-(4-aminophenyl)-2-pyrrolidinemethanols, as primary intermediates, in addition to other conventionally-used additives and components.

BACKGROUND OF THE INVENTION

[0002] Oxidative hair dye colorants are essential elements in hair dyeing preparations for the permanent dyeing of human hair. The hair dyeing process is achieved, in general, by the reaction of certain developing compounds with certain coupling compounds in the presence of a suitable oxidizing agent or compound, such as hydrogen peroxide.

[0003] When oxidation dyes, such as those comprising primary intermediates and couplers are used in the dyeing of human hair, the procedure may involve the use of a two part system. In general, one part can be a lotion formulation which contains a variety of ingredients, including oxidation dye precursors (i.e., primary intermediates and coupling agents). The other part is a developer formulation containing a suitable oxidizing agent, e.g., hydrogen peroxide. Immediately prior to application to the hair, the two parts are mixed to form a thickened liquid solution, for example, a lotion or a gel. As a consequence of the oxidizing properties of the oxidizing agent, some of the natural melanin pigment of the hair may be bleached. The precursors in the thickened solution (e.g. lotion or gel) penetrate into the hair where the primary intermediates are oxidized and react with the coupling agents to produce the desired color. Such systems generally contain a proportion of organic solvents and surfactants and contain relatively high levels of dye precursors to produce the desired color.

[0004] In order for procedures using permanent oxidative dyes to work properly, a number of parameters and conditions are important to consider in the use of these dyes in admixture with couplers in hair color preparations for human hair. Among these are the final color and color intensity that are produced after application to the subject's hair; the wash fastness and the light fastness of the resulting dye; the resistance of the dye to perspiration; the resistance of the dye to various hair treatments, such as permanent wave, straightening, shampooing, conditioning and rubbing. In addition, the dye must have virtually no allergenicity or dermal or systemic toxicity.

[0005] p-Phenylenediamine plays a very important role in oxidative hair coloring because a majority of shades are obtained with dyes based on this dye. However, the hair coloring industry is searching for p-phenylenediamine alternatives possessing better allergenic profiles than p-phenylenediamine. As part of the solution, GB 2,239,265A describes that 2-(2-hydroxyethyl)-p-phenylenediamine can be used as a potential replacement for p-phenylenediamine. U.S. patent 5,538,516 also describes the use of 2-(hydroxyalkoxy)-p-phenylenediamine as a p-phenylenediamine substitute. European patent 634,164 lists more than 40 compounds of p-phenylenediamine derivatives. Among them, N-4-aminophenylmorpholine and N-4-aminophenylpiperidine were described. Morpholine and piperidine derivatives are examples of heterocyclic compounds. No example of pyrrolidine was found in the list. However, these compounds are not good primary intermediates because the energy required for the formation of the oxidized forms of the piperidine and morpholine derivative is higher than for the open chain compound such as 4-amino-N,N-diethylaniline (R.L. Bent, J.C. Dessloch, F.C. Duennbier, D.W. Fassett, D.B. Glass, T.H. James, D.B. Julian, W.R. Ruby, J.M. Snell, J.H. Sterner, J.R. Thirtle, P.W. Vittum and A. Weissberger, J. Am. Chem. Soc. 1951, 73, 1300). It was also reported in this paper that formation of the oxidized compound is even more favorable for 1-(4-aminophenyl) pyrrolidine than for the 4-aminodialkylanilines with open alkyl groups such as 4-amino-N,N-diethylaniline. Although 1-(4-aminophenyl)pyrrolidine has a good potential as a primary intermediate, it was found to exhibit a strong allergenic activity. Because of adverse biological activity and various limitations, there is need for developing pyrrolidine derivatives which can be used as p-phenylenediamine alternatives.

[0006] In R.L. Bent, J.C. Dessloch, F.C. Duennbier, D.W. Fassett, D.B. Glass, T.H. James, D.B. Julian, W.R. Ruby, J.M. Snell, J.H. Sterner, J.R. Thirtle, P.W. Vitt, A. Weissberger, Chemical constitution, Electrochemical, Photographic and allergenic properties of p-amino-N-dialkylanilines, J. Am. Chem. Soc. 1951, 73, 1300, a total of 60 of 4-amino-N, N-dialkylanilines were studied on allergenic properties. 1-(4-Aminophenyl)pyrrolidine was one of them mentioned in the studies.

Piniella et al., "Synthesis, structure and second-order nonlinear optical properties of highly functionalised 6-aminopentafulvenes" J. Mater. Chem. 1998; Vol. 8(3), March 1998; pp 619-627 discloses N-(p-aminophenyl)prolinol and its mixture with a trihydroxypentafulvene at room temperature.

US 5,728,034 relates to N-(4-aminophenyl)pyrrolidine derivatives for use in photographic development.

SUMMARY OF THE INVENTION


[0007] It is an object of the invention to provide oxidative hair dye compositions and methods for the oxidative coloration of hair comprising compounds of the class of 1-(4-aminophenyl)-2-pyrrolidinemethanols, or cosmetically acceptable salts thereof, as primary intermediates in compositions comprising such primary dye intermediates as well as coupling agents, oxidizing agents and other adjuvant substances. In accordance with a particular aspect of the invention 1-(4-aminophenyl)-2-pyrrolidinemethanol is preferred.

[0008] It is a further object of the invention to use 1-(4-aminophenyl)-2-pyrrolidinemethanol compounds that serve as effective and high-performance primary intermediates in oxidative hair dyes. The 1-(4-aminophenyl)-2-pyrrolidinemethanol compounds are capable of replacing, in whole or in part, p-phenylenediamine (PPD), a known allergen and a sensitizer, and can be employed with couplers as color modifiers in such hair dye formulations.

[0009] Yet another object of the invention is to use a newly-discovered primary dye intermediate, 1-(4-aminophenyl)-2-pyrrolidinemethanol, or a cosmetically acceptable salt thereof, which provides intense color, when used in oxidative hair dye compositions in combination with coupling agents, oxidizing agents and other hair dye additives and/or adjuvants conventionally used in hair dye formulations for the dyeing of hair.


[0010] Further objects and advantages afforded by the invention will be apparent from the detailed description hereinbelow.

[0011] Accordingly, in a first aspect the present invention provides use of a compound having the formula I:




wherein R₁ is hydrogen, C₁-C₆ alkyl and C₁-C₅ mono or polyhydroxyalkyl, and R₂ is H or OH, or a cosmetically acceptable salt thereof, as a primary dye intermediate in the manufacture of an oxidative hair dye product.

[0012] In a further aspect, the present invention provides a composition for the oxidative colouring of hair comprising at least one primary dye intermediate; at least one coupling compound; and a cosmetically acceptable vehicle, wherein the at least one primary dye intermediate comprises a 1-(4-aminophenyl)-2-pyrrolidinemethanol, or a cosmetically acceptable salt thereof, having the formula I:



wherein R_1 is hydrogen, C_1 - C_6 alkyl and C_1 - C_5 mono or polyhydroxyalkyl, and R_2 is H or OH,
and wherein said composition further includes one or more additional dye components selected from the group
consisting of: p-phenylenediamines and cosmetically acceptable derivatives thereof; p-aminophenols and cosmetically
acceptable derivatives thereof; ortho developers and cosmetically acceptable derivatives thereof; phenols and cos-
metically acceptable derivatives thereof; resorcinols and cosmetically acceptable derivatives thereof; m-phenylenedi-
amines and cosmetically acceptable derivatives thereof; m-aminophenols and cosmetically acceptable derivatives
thereof; and heterocyclic derivatives.


[0013] In a further aspect, the present invention provides a composition for the oxidative colouring of hair comprising
at least one primary dye intermediate; at least one coupling compound; and a cosmetically acceptable vehicle, wherein
the at least one primary dye intermediate comprises a 1-(4-aminophenyl)-2-pyrrolidinemethanol, or a cosmetically
acceptable salt thereof, having the formula I:



wherein R_1 is hydrogen, C_1 - C_6 alkyl and C_1 - C_5 mono or polyhydroxyalkyl, and R_2 is H or OH,
and wherein the composition further includes an oxidising agent.

DETAILED DESCRIPTION OF THE INVENTION

[0014] The present invention relates to primary intermediates for use in oxidative hair dye compositions. The com-
positions of the present invention include, as primary dye intermediates, one or more components from the chemical
class of substituted or unsubstituted 1-(4-aminophenyl)-2-pyrrolidinemethanol compounds and cosmetically accepta-
ble salts thereof. Nonlimiting examples of derivatives and salts of the 1-(4-aminophenyl)-2-pyrrolidinemethanol com-
pounds include sulfates, hydrochlorides, phosphates and the like with sulfates being preferred. The 1-(4-aminophenyl)
pyrrolidine compounds for use in the present invention are generally represented by the following



chemical formula I:


wherein R_1 is a hydrogen; C_1 - C_6 alkyl, preferably C_1 - C_4 alkyl; or C_1 - C_5 mono- or polyhydroxyalkyl, and R_2 is hydrogen or hydroxyl group. Preferred are unsubstituted 1-(4-aminophenyl)-2-pyrrolidinemethanol (e.g., formula I, wherein R_1 and R_2 are hydrogen), as well as substituted 1-(4-aminophenyl)-2-pyrrolidinemethanols (e.g., formula I, wherein R_1 is methyl or ethyl and R_2 is hydrogen), and 1-(4-aminophenyl)-4-hydroxy-2-pyrrolidinemethanol (e.g., formula I, where R_1 is hydrogen and R_2 is hydroxyl group).

[0015] The compositions of the present invention contain as a primary dye intermediate one or more substituted or unsubstituted 1-(4-aminophenyl)-2-pyrrolidinemethanol compounds, or a cosmetically acceptable salt thereof, which may be present in admixture with one or more other oxidation dye precursor(s), for example, primary dye intermediate or conventional coupling component(s), in addition to other hair dye component ingredients, additives, or adjuvants typically used in oxidative hair dye formulations as described herein. The 1-(4-aminophenyl)-2-pyrrolidinemethanol, alone or in combination with one or more other dye intermediates and coupling agents produce novel dyestuffs which provide intense coloration to hair. The 1-(4-aminophenyl)-2-pyrrolidinemethanol compounds discussed herein achieve a deep coloration to hair and impart unique properties to dye substances used for oxidative hair coloring.

[0016] The compounds of the class of 1-(4-aminophenyl)-2-pyrrolidinemethanols, or cosmetically acceptable salts thereof, are distinct from hair dyeing compounds disclosed in the prior art and offer newly-discovered and advantageous hair coloring properties.

[0017] It will be appreciated by those having skill in the art that the compositions, methods and uses of the present invention are appropriate for the dyeing of keratinous fibers, including the hair fibers of animals and humans, with particular application to the oxidative coloring of human hair.

[0018] It has been found surprisingly that dyes produced by coupling the compound 1 or 1a with conventional couplers are different from those derived from coupling 1-(4-aminophenyl)pyrrolidine 2 or N,N-bis(2-hydroxyethyl)-p-phenylenediamine 3 with the same couplers in spite of the fact that the compounds are structurally similar to each other.



[0019] For example, 1-(4-aminophenyl)-2-pyrrolidinemethanol (**1**) couples with 3-methyl-1-phenyl-2-pyrazolin-5-one to produce a dye which colors piedmont hair intense dark violet. The compound of formula **1a** also produces violet when coupled with 3-methyl-1-phenyl-2-pyrazolin-5-one. In contrast, 1-(4-aminophenyl)pyrrolidine (**2**) couples with 3-methyl-1-phenyl-2-pyrazolin-5-one to produce a dye which imparts a reddish violet coloration to piedmont hair. N,N-bis(2-hydroxyethyl)-p-phenylenediamine (**3**) imparts blue violet coloration to piedmont hair. These visual differences are well manifested in the CIE L*a*b* values (compare entries 1, 7 and 12 in Table 2). In the chromaticity diagram, -b* is the blue direction.

[0020] 2,4-Diaminophenoxyethanol couples with the novel primary intermediate of the present invention to produce a dye which colors piedmont hair intense blue. The blue color obtained is different from that derived from 1-(4-aminophenyl)pyrrolidine **2** and 2,4-diaminophenoxyethanol. b* Value of the former is -12.98, while the latter b* is -9.36 (Table 2). The dye on gray hair exhibits excellent wash and light fastness (Table 5).

[0021] 2-Methyl-1-naphthol couples with **1** to impart a blue color on piedmont hair. Again, the compound **1** provides a b* value -20.3 which is substantially lower than the counterpart b* value -13.07 obtained with the compound **2**.

[0022] 3-Aminophenol couples with **1** to produce a dye which colors hair dark black, while compound **2** colors hair neutral black. The color on gray hair exhibits excellent wash and light fastness properties (Table 4).

[0023] Overall wash and light fastness of **1** is better than those of N,N-bis(2-hydroxyethyl)-p-phenylenediamine **3** when coupled with m-aminophenol or 2,4-diaminophenoxyethanol as shown in Table 4 and 5. The smaller ΔE is, the better wash and light fastness is.

[0024] Violet color obtained from **1** and 3-methyl-1-phenyl-2-pyrazolin-5-one is very similar to that produced from p-phenylenediamine and 2-methyl-5-aminophenol (see entries 1 and 13 in Table 2). This observation may allow one to formulate a red shade without relying on p-phenylenediamine. The red shade is generally produced by a combination of p-phenylenediamine, p-aminophenol and 2-methyl-5-aminophenol.

[0025] As described above, in addition to at least one of the novel component dye molecules encompassed by the present invention, the hair dyeing compositions described herein may also contain at least one other known and usual oxidation dye precursor (i.e., used as primary dye intermediates and/or couplers), as well as conventional direct colorants and dyes in admixture, should these substances be necessary or desired for the development and production of certain color nuances and tints.

[0026] Illustrative component dye ingredients that are conventionally admixed and employed as constituents of customary hair dye formulations and that can be considered suitable for use in the compositions of the present invention are set forth hereinbelow. As one particular example, p-phenylenediamine, used in oxidative hair coloring formulations, may conveniently be used in admixture with the novel primary intermediate 1-(4-aminophenyl)-2-pyrrolidinemethanol compounds in the compositions of the present invention.

[0027] Included among the suitable dye components that may be considered for use as primary intermediates and/or couplers in the dye compositions of the present invention are the following: p-phenylenediamine derivatives such as: p-toluenediamine; p-phenylenediamine; 2-chloro-p-phenylenediamine; N-phenyl-p-phenylenediamine; N-2-methoxyethyl-p-phenylenediamine; N,N-bis-(hydroxyethyl)-p-phenylenediamine; 2-hydroxymethyl-p-phenylenediamine; 2-hydroxyethyl-p-phenylenediamine; 4,4'-diaminodiphenylamine; 2,6-dimethyl-p-phenylenediamine; 2-isopropyl-p-phenylenediamine; N-(2-hydroxypropyl)-p-phenylenediamine; 2-propyl-p-phenylenediamine; 1,3-N, N-bis-(2-hydroxyethyl)-N, N-bis (4-aminophenyl)-2-propanol; and 2-methyl-4-dimethylaminoaniline, or combinations thereof.

[0028] Preferred p-phenylenediamine derivatives include: p-toluenediamine; p-phenylenediamine; N-2-methoxyethyl-p-phenylenediamine; N,N-bis(hydroxyethyl)-p-phenylenediamine; and 2-hydroxyethyl-p-phenylenediamine.

[0029] p-Aminophenol derivatives include: p-aminophenol; p-methylaminophenol; 3-methyl-p-aminophenol; 2-hydroxymethyl-p-aminophenol; 2-methyl-p-aminophenol; 2-(2-hydroxyethylaminomethyl)-p-aminophenol; 2-methoxymethyl-p-aminophenol; and 5-aminosalicylic acid, or combinations thereof.

[0030] Preferred p-aminophenol derivatives include: p-aminophenol; p-methylaminophenol; 3-methyl-p-aminophenol; 2-methyl-p-aminophenol; 2-(2-hydroxyethylaminomethyl)-p-aminophenol; 2-methoxymethyl-p-aminophenol; and 5-aminosalicylic acid.

[0031] Ortho developers include: catechol; pyrogallol; o-aminophenol; 2,4-diaminophenol; 2,4,5-trihydroxytoluene; 1,2,4-trihydroxybenzene; 2-ethylamino-p-cresol; 2,3-dihydroxynaphthalene; 5-methyl-o-aminophenol; 6-methyl-o-aminophenol; and 2-amino-5-acetaminophenol, or combinations thereof.

[0032] Preferred ortho developers include: o-aminophenol; 2,4-diaminophenol; 2,4,5-trihydroxytoluene; 1,2,4-trihydroxybenzene; 2-ethylamino-p-cresol; 5-methyl-o-aminophenol; 6-methyl-o-aminophenol; and 2-amino-5-acetaminophenol.

[0033] Phenols, naphthols and resorcinol derivatives include: 2-methyl-1-naphthol; 1-acetoxy-2-methylnaphthalene; 1,7-dihydroxynaphthalene; resorcinol; 4-chlororesorcinol; 1-naphthol; 1,5-dihydroxynaphthalene; 2,7-dihydroxynaphthalene; 2-methylresorcinol; 1-hydroxy-6-aminonaphthalene-3-sulfonic acid; thymol (2-isopropyl-5-methylphenol); 1,5-dihydroxy-1,2,3,4-tetrahydronaphthalene; 2-chlororesorcinol; 2,3-dihydroxy-1,4-naphthoquinone; and 1-naphthol-4-sulfonic acid, or combinations thereof.

[0034] Preferred phenols, naphthols and resorcinol derivatives include: 2-methyl-1-naphthol; 1-acetoxy-2-methyl-naphthalene; 1,7-dihydroxynaphthalene; resorcinol; 4-chlororesorcinol; 1-naphthol; 1,5-dihydroxynaphthalene; 2,7-dihydroxynaphthalene; 2-methylresorcinol; thymol (2-isopropyl-5-methylphenol); and 2,3-dihydroxy-1, 4-naphthoquinone.

5 [0035] m-Phenylenediamines include: m-phenylenediamine; 2-(2,4 diaminophenoxy)ethanol; N,N-bis(hydroxyethyl)-m-phenylenediamine; 2,6-diaminotoluene; N,N-bis(hydroxyethyl)-2,4-diaminophenetole; bis(2,4-diaminophenoxy)-1,3-propane; 1-hydroxyethyl-2, 4-diaminobenzene; 2-amino-4 hydroxyethylaminoanisole; aminoethoxy-2,4-diaminobenzene; 2,4-diaminophenoxyacetic acid; 4,6-bis(hydroxyethoxy)-m-phenylenediamine; 2,4-diamino-5-methylphenetole; 2,4-diamino-5-hydroxyethoxytoluene; 2,4-dimethoxy 1,3-diaminobenzene; and 2,6-bis(hydroxyethylamino)toluene, or combinations thereof.

10 [0036] Preferred m-phenylenediamines include: m-phenylenediamine; 2,4-diaminophenoxyethanol; bis-(2,4-diaminophenoxy)-1,3-propane; 1-hydroxyethyl-2,4-diaminobenzene; 2-amino-4-hydroxyethylaminoanisole; 4,6-bis(hydroxyethoxy)-m-phenylenediamine; 2,4-diamino-5-methylphenetole; 2,4-diamino-5-hydroxyethoxytoluene; 2,4-dimethoxy-1,3-diaminobenzene; and 2,6-bis(hydroxyethylamino)toluene.

15 [0037] m-Aminophenols include: m-aminophenol; 2-hydroxy-4-carbamoylmethylaminotoluene; m-carbamoylmethylaminophenol; 6-hydroxybenzomorpholine; 2-hydroxy-4-aminotoluene; 2-hydroxy-4-hydroxyethylaminotoluene; 4,6-dichloro-m-aminophenol; 2-methyl-m-aminophenol; 2-chloro-6-methyl-m-aminophenol; 2-hydroxyethoxy-5-aminophenol; 2-chloro-5-trifluoroethylaminophenol; 4-chloro-6-methyl-m-aminophenol; N-cyclopentyl-3-aminophenol; N-hydroxyethyl-4-methoxy-2-methyl-m-aminophenol and 5-amino-4-methoxy-2-methylphenol, or combinations thereof.

20 [0038] Preferred m-aminophenols include: m-aminophenol; 6-hydroxybenzomorpholine; 2-hydroxy-4-aminotoluene; 2-hydroxy-4-hydroxyethylaminotoluene; 4,6-dichloro-m-aminophenol; 2-methyl-m-aminophenol; 2-chloro-6-methyl-m-aminophenol; 4-chloro-6-methyl-m-aminophenol; N-cyclopentyl-3-aminophenol; N-hydroxyethyl-4-methoxy-2-methyl-m-aminophenol and 5-amino-4-methoxy-2-methylphenol.

25 [0039] Heterocyclic derivatives include: 2-dimethylamino-5-aminopyridine; 2,4,5,6-tetra-aminopyrimidine; 4,5-diamino-1-methylpyrazole; 1-phenyl-3-methyl-5-pyrazolone; 6-methoxy-8-aminoquinoline; 2,6-dihydroxy-4-methylpyridine; 5-hydroxy-1,4-benzodioxane; 3,4-methylenedioxyphenol; 4-hydroxyethylamino-1,2-methylenedioxybenzene; 2,6-dihydroxy-3,4-dimethylpyridine; 5-chloro-2,3-dihydroxypyridine; 3,5-diamino-2,6-dimethoxypyridine; 2-hydroxyethylamino-6-methoxy-3-aminopyridine; 3,4-methylenedioxyaniline; 2,6-bis-hydroxyethoxy-3,5-diaminopyridine; 4-hydroxyindole; 3-amino-5-hydroxy-2,6-dimethoxypyridine; 5,6-dihydroxyindole; 7-hydroxyindole; 5-hydroxyindole; 2-bromo-4,5-methylenedioxyphenol; 6-hydroxyindole; 3-amino-2-methylamino-6-methoxypyridine; 2-amino-3-hydroxypyridine; 2,6-diaminopyridine; 5-(3,5-diamino-2-pyridyloxy)-1,3-dihydroxypentane; 3-(3,5-diamino-2-pyridyloxy)-2-hydroxypropanol and 4-hydroxy-2,5,6-triaminopyrimidine, or combinations thereof.

30 [0040] Preferred heterocyclic derivatives include: 4,5-diamino-1-methyl-pyrazole; 2-dimethylamino-5-aminopyridine; 2,4,5,6-tetra-aminopyrimidine; 1-phenyl-3-methyl-5-pyrazolone; 3,4-methylenedioxyphenol; 4-hydroxyethylamino-1,2-methylenedioxybenzene; 2,6-dihydroxy-3,4-dimethylpyridine; 5-chloro-2,3-dihydroxypyridine; 3,5-diamino-2,6-dimethoxypyridine; 2-hydroxyethylamino-6-methoxy-3-aminopyridine; 3,4-methylenedioxyaniline; 4-hydroxyindole; 5,6-dihydroxyindole; 7-hydroxyindole; 5-hydroxyindole; 2-bromo-4,5-methylenedioxyphenol; 6-hydroxyindole; 3-amino-2-methylamino-6-methoxypyridine; 2-amino-3-hydroxypyridine; 2,6-diaminopyridine; 5-(3,5-diamino-2-pyridyloxy)-1,3-dihydroxypentane; 3-(3,5-diamino-2-pyridyloxy)-2-hydroxypropanol and 4-hydroxy-2,5,6-triaminopyrimidine.

40 [0041] Particularly preferred couplers for use in combination with the compounds of the present invention are resorcinol, m-aminophenol, 2,4-diaminophenoxyethanol and 2-methyl-1-naphthol, which may be used alone or in combinations of two or more. The most preferred compositions of the present invention comprise 1-(4-aminophenyl)-2-pyrrolidinemethanol 1 and/or N-(4-aminophenyl)-4-hydroxy-2-pyrrolidinemethanol 1a in combination with a coupler or coupler system selected from:

- 45
- (a) m-aminophenol;
 - (b) resorcinol;
 - (c) m-aminophenol and resorcinol;
 - 50 (d) resorcinol and 2,4-diaminophenoxyethanol;
 - (e) m-aminophenol and 2-methyl-1-naphthol;
 - (f) 2,4-diaminophenoxyethanol and m-aminophenol, and
 - (g) resorcinol and 2-methyl-1-naphthol.

55 These combinations produce in combination with Compounds 1 and 1a neutral black, dark black and blue black shades.

[0042] The additional dye compounds, e.g., couplers, should be present in the hair dyeing compositions of the present invention in an amount of approximately 0.01 to 10%, by weight, preferably approximately 0.1 to 5%, by weight, based on the total weight of the composition. The total quantity of oxidative colorant, comprising primary intermediate(s) and

coupler(s), will suitably amount to approximately 0.1 to 10%, by weight, and preferably, approximately 0.5 to 5% by weight, based on the total weight of the composition.

[0043] Unless indicated otherwise, as used herein, reagent or component amounts are in % by weight (w/w), based on the total weight of the composition.

5 [0044] In the compositions of the present invention, the coupling component is generally used in approximately equimolar amounts relative to the primary intermediate (developing component). However, it will be appreciated that the primary intermediate in relation to the coupler, may be present either in increased or decreased amounts depending upon the formulation and the desired color, intensity or effect. In general terms, the primary intermediate and the coupling component, or cosmetically acceptable salts thereof, will be present in tinctorially effective amounts for the coloring of a hair fiber.

10 [0045] The hair dye preparations of the present invention may be formulated into cosmetic preparations such as a solution, cream, lotion, gel or emulsion. Also, in accordance with the invention, the compositions may represent a mixture of the coloring components (i.e., dye intermediate and coupling agent) with other components commonly associated with the formulation of solutions, creams, lotions, gels or emulsions, and the like. For example, components 15 such as wetting agents or emulsifying agents from the categories of anionic or nonionic surfactants, such as sulfates of fatty alcohols, alkanolamides of fatty alcohols, alkyl sulfonates, alkylbenzene sulfonates, oxyethylated fatty alcohols, oxyethylated nonylphenols; furthermore thickeners such as fatty alcohols, starch, cellulose derivatives, paraffin oil and fatty acids, as well as hair-care substances such as lanolin derivatives, cholesterol and pantothenic acid, may be formulated into the compositions of the invention.

20 [0046] As an example, if formulated as a lotion, the compositions of the invention may contain organic solvents to assist in dissolving the dye precursors. Accordingly, the organic solvent content of the lotion may be from 0% to about 20%, preferably, about 1% to about 15%. Typically useful solvents include alcohols containing up to three carbon atoms such as ethanol and isopropanol, polyhydroxy alcohols such as propylene or hexylene glycol and lower alkyl ethers thereof, such as ethoxy ethers.

25 [0047] In addition, the hair dyeing compositions in accordance with the present invention may optionally contain conventionally-used adjuvants and cosmetic additives, or mixtures thereof, to achieve the final formulations. Examples of such additives include, but are not limited to, anti-oxidants, e.g., ascorbic acid, erythorbic acid, or sodium sulfite, to inhibit premature oxidizing; fragrances and/or perfume oils; chelating agents; emulsifiers; coloring agents; thickeners; organic solvents; opacifying agents; dispersing agents; sequestering agents; hair-care substances; humectants; and antimicrobials, and others. The list of optional ingredients is not intended as limiting. Other suitable adjuvants for inclusion in the hair dye compositions of the invention are disclosed, for example, in Zviak, *The Science of Hair Care* (1986) and in Balsam and Sagarin, *Cosmetics: Science and Technology*, Vol. 2, Second Edition, (1972).

30 [0048] Thickeners that may be used in the compositions of the present invention include a variety of fatty acid soaps and associative polymeric thickeners. The fatty acid soaps are alkaline metal salts or alkanolamine salts of fatty acids with C₁₀-C₁₆ alkyl side chains. The preferred fatty acids include oleic acid, myristic acid and lauric acid, which are generally present in the compositions of the invention at about 0.5% to about 20%, preferably about 1% to about 10%. Associative thickeners are polymers that can thicken solutions at low concentrations. Among the associative thickeners that are useful in the compositions of the present invention are acrylates copolymer (sold by Rohm and Haas under the tradename Aculyn-33), cetareth-20 acrylates/steareth-20 methacrylate copolymer (sold by Rohm and Haas under the Trade Name Aculyn22), acrylates/steareth-20 itaconate copolymer and acrylates/ceteth-20 itaconate copolymer. Another class of associative thickeners useful in the compositions of the present invention include the copolymers of polyurethane and polyethylene glycol or polyether urethanes. One such material is sold by Rohm and Haas under the tradename Aculyn-44. The associative polymeric thickeners are generally present in the compositions of the invention at about 0.1 % to about 10%, preferably, about 0.5% to about 5%.

45 [0049] The oxidative coupling, i.e., the development of the dye, to produce the final color product on the hair, can, in principle, be performed with atmospheric oxygen. However, chemical oxidizing agents are suitably and preferably used. Although other oxidizing agents can be employed, hydrogen peroxide is a preferred oxidizing compound for use as a developer with the primary intermediate and coupler dye precursors of the invention. The concentration of hydrogen peroxide in the developer may be from about 1% to about 15%, preferably, from about 3% to about 12%. Other suitable oxidizing agents include, for example, urea peroxide, melamine peroxide, perborates and percarbonates such as sodium perborate or percarbonate. The amounts of such oxidizing agents can be routinely determined by one having skill in the art, without requiring any inventive skill.

50 [0050] The compositions of the invention may include a typical anionic, cationic, nonionic or amphoteric surfactant. The anionic surfactants include the variety of alkyl sulfates, alkylether sulfates, alkyl sulfonates, alkyl sulfosuccinates and N-acyl sarcosinates. The commonly used anionic surfactants are sodium and ammonium lauryl sulfates, sodium and ammonium laureth sulfate and alpha olefin sulfonates. Anionic surfactants are generally present in the compositions of the present invention at about 0.1 % to about 15%, preferably, about 0.5% to about 10%.

55 [0051] The nonionic surfactants that can be used in the present invention include the wide variety of ethoxylated

alcohols, nonoxynols, alkanolamides, alkyl stearates, alkyl palmitates and alkylpolyglucosides. Examples of the commonly used nonionic surfactants are cetyl alcohol, stearyl alcohol, oleyl alcohol; the various types of ethoxylated alkylphenols; lauroyl diethanolamide; lauroyl monoethanolamide; isopropyl palmitate, isopropyl stearate and decylpolyglucoside. Nonionic surfactants are generally present in the compositions of the present invention at about 0.1% to about 15%, preferably, about 0.5% to about 10%.

[0052] The compositions in accordance with the present invention may also contain one or more quaternary ammonium compounds that provide hair conditioning effects. The quaternary ammonium compounds can be monomeric or polymeric quaternary ammonium compounds. Nonlimiting examples of such compounds include cetyltrimonium chloride, stearyl trimonium chloride, benzalkonium chloride, behentrimonium chloride and a variety of polyquaterniums. The quaternary ammonium compounds are generally present in the compositions of the present invention at about 0.1% to about 10%, preferably, about 0.5% to about 5%.

[0053] Amphoteric surfactants that can be incorporated in the compositions of the present invention belong to the class of surface active chemicals that possess a positive and a negative charge in the same molecule and behave as a cation, an anion, or both, depending upon the pH of the medium and the nature of the amphoteric molecule. In general, the positive charge is located on a nitrogen, while the negative charge is carried by a carboxyl or sulfonate group. There are a large number of amphoteric surfactants that are suitable for use in the present invention, including, for example, the well-known betaines, sultaines, glycinate and propionates.

[0054] The selection of the amphoteric surfactant or mixture of surfactants for use in the present compositions and methods is not critical. The surfactant may be selected from among those suggested above, or from any of a number of other known amphoteric surfactants. The amount of amphoteric surfactant in the compositions of the present invention is normally from about 0.5% to about 15%, preferably, about 2% to about 10%.

[0055] Depending on the final formulated preparation, the compositions in accordance with invention may be weakly acidic, neutral or alkaline. In particular, the pH of the prepared compositions can range from about 5 to about 11. Preferred is a pH range of about 8 to 10. Any of a wide variety of alkaline reagents can be used to adjust the pH of the hair coloring compositions. Such alkaline reagents include ammonium hydroxide, potassium or calcium hydroxide, sodium or potassium carbonate, sodium phosphate, sodium silicate, guanidine hydroxide, or any one of the alkylamines or alkanolamines, for example, ethylamine, triethylamine, trihydroxymethylamine, ethanolamine, diethanolamine, aminomethyl propanol, aminomethyl propanediol and the like. The preferred alkaline reagents are ammonium hydroxide, sodium carbonate and ethanolamine. With the reagents listed above, the selected pH will generally be achieved if the composition contains an alkaline agent in an amount from about 0.1% to about 15%, preferably, about 0.5% to about 5%.

[0056] The application of the dyeing components is carried out by methods familiar to those in the art, for example, by mixing the hair dyeing preparation with an oxidant shortly before use, or at the time of applying the mixture onto the hair. On the hair, the compositions form a stable formulation with enough consistency and body to remain on the hair without dripping or running during the complete coloring period. The primary intermediate and coupler, i.e. the dye precursors, diffuse rapidly into the hair together with the oxidizing agent, or developer. The dyes form within the hair fiber, and since they are large molecules, remain in the hair so that the color change is permanent. The term "permanent" means the dye does not readily wash out of the hair with ordinary shampoos. At the end of coloring application (e.g., approximately 5 to 45 minutes, preferably, approximately 10 to 30 minutes), the composition is washed from the hair with an ordinary water rinse followed by a shampoo. The application temperature is in the range of about 15°C to 50°C.

[0057] The 1-(4-aminophenyl) pyrrolidine methanols and cosmetically-acceptable derivatives and salts thereof, are soluble in water and have a long shelf life, particularly as constituents of the hair dyeing preparations in accordance with the invention. These primary intermediates, of which 1-(4-aminophenyl)-2-pyrrolidinemethanol is preferred, should be present in the hair dyeing preparations in an amount of approximately 0.1% to approximately 10%, preferably, approximately 0.1% to approximately 5%. The total quantity of oxidative colorant, consisting of primary intermediate(s) and coupling component(s) will suitably amount to approximately 0.1% to approximately 20%, and preferably, approximately 0.5% to approximately 15% of the composition.

[0058] The hair dyeing compounds in accordance with the invention will offer a wide range of varying color tints depending upon the type and composition of the colorant constituents. The color tints are distinguished herein by their particular intense and lasting color. The superior coloring properties of the hair dyeing compositions of the present invention are further evidenced by allowing grayed hair that has not been subjected to prior chemical damage to be covered without problems and with a depth and covering strength that, prior to the discovery of the primary intermediates of the present invention, had only been attained using the conventional PPD.

[0059] The compositions of this invention may be separately provided in a kit or packaged form ready for mixing by the user, either professional or consumer, to initiate the dyeing process. The kit provided in accordance with this invention comprises containers for housing the developer and the dye precursors. In the most convenient form, there will be two containers, one containing the primary dye intermediate and coupler, e.g., as a lotion; the other containing the oxidizing agent, also called the developer or developing agent.

[0060] The method of the invention comprises applying the mixture to the hair to be colored and allowing it to remain

in contact with the hair until the desired hair color has been attained, after which time the composition is removed from the hair as described above.


EXAMPLES

[0061] The examples as set forth herein are meant to exemplify the various aspects of carrying out the invention and are not intended to limit the invention in any way.

Example 1. The synthesis of compound 1

[0062] 1-Fluoro-4-nitrobenzene 4 (13.3 g, 94 mmole) is treated with (s)-(+)-2-pyrrolidinemethanol 5 (9.5 g, 94 mmole) and potassium carbonate (15.6 g, 113 mmole) in DMF (188 ml, 0.5M solution) at 60°C for 6 hours. The reaction is cooled to room temperature. The solvent is evaporated under vacuum and the residue is taken up into dichloromethane (500 ml). The organic phase is washed with saturated NaCl solution (250 ml). The phases are separated and the organic phase is dried over MgSO₄, filtered, and evaporated under vacuum to leave an orange oil. The crude material is then crystallized from ethyl acetate/hexane to produce compound 6 (18.3 g, 88% yield) as a bright yellow solid: mp 118-120°C; ¹H NMR (300 MHz, DMSO-d₆) δ 2.06 (m, 4H), 3.31 (m, 2H), 3.55 (d, 2H, J = 9.0 Hz), 3.98 (s, 1H), 4.97 (t, 1H, J = 6.0 Hz), 6.77 (d, 2H, J = 8.7 Hz), 8.12 (d, 2H, J = 8.7 Hz); m/z 222 (M⁺).

[0063] Compound 6 (13.3 g, 60 mmole) is then hydrogenated at 60 psi of hydrogen in the presence of 10% Pd on carbon (1.3 g) in ethanol (222 ml, 0.27M solution) for 2 hours at room temperature. The solution is then filtered through a pad of Celite, evaporated, and the crude material is recrystallized from ethyl acetate/hexane to give compound 1 (10.1 g, 88% yield) as a light grey solid: mp 94-95°C; ¹H NMR (300 MHz, DMSO-d₆) δ 1.86 (m, 4H), 2.84 (m, 1H), 3.11 (m, 1H), 3.29 (m, 3H), 4.29 (bs, 2H), 4.66 (bs, 1H), 6.38 (d, 2H, J = 8.1 Hz), 6.48 (d, 2H, J = 8.1 Hz); m/z 192 (m⁺).




Example 2. The synthesis of compound 1a

[0064] 1-Fluoro-4-nitrobenzene 4 (42.3 g, 0.3 mole) is treated with trans-4-hydroxy-L-proline 7 (39.3 g, 0.3 mole) in the presence of potassium fluoride (17.4 g, 0.3 mole) and 18-Crown-6 (6.1g, 30.0 mmole) in DMSO (300 ml) at 60 °C for 18 hours. The reaction is cooled to room temperature and poured into an ice chilled saturated NaCl solution (500

ml). The resulting aqueous solution is acidified to pH about 3 with aqueous 10% HCl. The aqueous phase is extracted with EtOAc (3 x 500 ml). The organic phases are combined, dried over MgSO_4 , and the solvent is vacuum evaporated. The crude material is then recrystallized from EtOAc/hexane to afford compound **8** (71.1 g, 94% yield) as a dark yellow solid: mp 98-100°C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 2.18-2.28 (m, 2H), 3.32 (dd, 1H, $J = 10.0, 2.1$ Hz), 3.65 (dd, 1H, $J = 11.0, 5.0$ Hz), 4.43 (m, 2H), 5.31 (s, 1H), 6.62 (d, 2H, $J = 9.0$ Hz), 8.12 (d, 2H, $J = 9.0$ Hz), 13.10 (bs, 1H); m/z 252 (M^+).

[0065] Compound **8** (31.5 g, 125 mmole) is then reduced with 1.0M BH_3 -THF complex (375 ml, 375 mmole) in THF (125 ml) at room temperature overnight (18 hrs). The excess BH_3 -THF complex is quenched with the slow addition of MeOH (100 ml). The solvent is then vacuum evaporated. The residue is taken up into EtOAc (500 ml) and an ice chilled saturated NaCl solution (500 ml). The aqueous phase is extracted with EtOAc (3 x 500ml). EtOH (50 ml) is added due to solubility problems. The organic phases are combined, dried over MgSO_4 , and the solvent is vacuum evaporated. The crude material is recrystallized from EtOAc/hexane to give compound **9** (13.0 g, 44% yield) as a yellow solid: mp 155-157°C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.96-2.03 (m, 1H), 2.56 (m, 1H), 3.18 (dd, 1H, $J = 10, 7.4$ Hz), 3.50 (t, 2H, $J = 5.1$ Hz), 3.68 (dd, 1H, $J = 11, 6$ Hz), 4.09 (m, 1H), 4.55 (m, 1H), 4.90 (t, 1H, $J = 5.7$ Hz), 5.14 (d, 1H, $J = 4.5$ Hz), 6.73 (d, 2H, $J = 9.3$ Hz), 8.08 (d, 2H, $J = 9.3$ Hz); m/z 238 (M^+).

[0066] Compound **9** (10.0 g, 42.0 mmole) is then hydrogenated at 60 psi of hydrogen in the presence of 10% Pd on carbon (1.5 g) in ethanol (200 ml, 0.2M solution) for 6 hours at room temperature. The solution is vacuum filtered through a pad of celite and the solvent is vacuum evaporated. The crude material is recrystallized from EtOAc/hexane to give compound **1a** (6.7 g, 75% yield) as a light tan solid: mp 130-132°C; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 1.74 (m, 1H), 2.02 (m, 1H), 2.67 (m, 1H), 3.09 (m, 1H), 3.39 (m, 1H), 3.49 (m, 1H), 4.20 (m, 2H), 4.31 (m, 1H), 4.55 (m, 1H), 4.81 (m, 1H), 6.28 (d, 2H, $J = 7.8$ Hz), 6.41 (d, 2H, $J = 7.8$ Hz); m/z 208 (M^+).



2. Dyeing procedure

[0067] The following compositions (1 or 1a or 2 or 3 or ppd and coupler: 5 mmole each) shown in table 1 are mixed with 100 g of 20 volume hydrogen peroxide. The resulting mixture is applied to piedmont hair or gray hair and permitted to remain in contact with hair for 30 minutes. Thus dyed hair is then shampooed and rinsed with water and dried. Table

2 shows the CIE L*a*b* values and colors obtained on piedmont hair.

Table 1.

Dyeing Composition	
	Composition(%)
Cocamidopropyl betaine	17
Ethanolamine	2
Oleic Acid	0.75
Citric Acid	0.1
Ammonium hydroxide	5.0
Behentrimonium chloride	0.5
Sodium sulfite	0.1
EDTA	0.1
Erythorbic acid	0.4
Primary intermediate (1, 1a, 2, 3 or ppd)	5 mmole
Coupler	5 mmole
Water	QS 100

Table 2.

Colors Obtained from Coupling of 1, 1a, 2, 3 or PPD with Various Couplers on Piedmont Hair						
Entry No.	PI	Coupler	L*	a*	b*	Color Description
1	1	3-Methyl-1-phenyl-2-pyrazolin-5-one	23.48	12.77	-1.09	intense violet
2	1a	3-Methyl-1-phenyl-2-pyrazolin-5-one	27.18	12.63	-4.83	Violet
3	1	2,4-Diaminophenoxyethanol	18.99	3.05	-12.98	Blue
4	1	2-Methyl-1-naphthol	21.33	6.16	-20.3	Blue
5	1	3-Aminophenol	21.06	0.22	-6.75	Black
6	1	2-Amino-3-hydroxypyridine	19.90	3.20	-3.90	Blue violet
7	2	3-Methyl-1-phenyl-2-pyrazolin-5-one	28.73	16.10	4.39	reddish violet
8		2,4-Diaminophenoxyethanol	18.27	2.47	-9.36	Blue
9		2-Methyl-1-naphthol	20.69	4.81	-13.07	Blue
10		3-Aminophenol	18.82	0.09	-3.71	Neutral black
11		2-Amino-3-hydroxypyridine	19.44	1.90	-1.62	Dark brown
12	3	3-Methyl-1-phenyl-2-pyrazolin-5-one	28.01	12.76	-4.49	Blue violet
13	PPD	2-Methyl-5-aminophenol	21.60	12.13	-0.79	Intense violet

Table 3.

Dyeing Composition of Reddish Brown Shade on Piedmont Hair	
	Composition(%)
Cocamidopropyl betaine	17.000
Ethanolamine	2.000
Oleic Acid	0.750
Citric Acid	0.100
Ammonium hydroxide	5.000
Behentrimonium chloride	0.500

Table 3. (continued)

Dyeing Composition of Reddish Brown Shade on Piedmont Hair	
	Composition(%)
Sodium sulfite	0.100
EDTA	0.100
Erythorbic acid	0.400
Primary intermediate 1 sulfate salt	0.725
p-Aminophenol	0.545
3-Methyl-1-phenyl-2-pyrazolin-5-one	0.440
2-Methyl-5-aminophenol	0.615
Water	QS 100.000
Shade	Reddish Brown

3. Wash and light fastness testing

[0068] Blended gray (50%) was used for the following experiments. For comparative purposes, N,N-bis(2-hydroxyethyl)-p-phenylenediamine 3 which is a common dye ingredient in commercial products was tested as a standard.

[0069] To test the degree of resistance to shampoo washing, the colored hair was immersed and shaken for 3 hours in a 10% Herbal Essences solution. Light fastness was tested by subjecting dyed hair to fade-o-meter for 72 hours.

Table 4.

Tristimulus Value of Shampoo and Light Fastness Studies with 3-aminophenol								
3-Aminophenol	PI	Before			After			ΔE
		L*	a*	b*	L*	a*	b*	
Wash 5 hr	1	16.83	-0.01	-2.18	17.19	0.04	-1.63	0.66
	2	15.21	0.06	-1.17	16.03	0.04	-1.20	0.82
	3	16.65	0.01	-1.89	18.33	-0.17	-1.53	1.73
Light 72 hour	1	16.83	-0.01	-2.18	18.17	-0.14	-0.15	2.44
	2	15.21	0.06	-1.17	15.58	-0.03	-0.19	1.05
	3	16.65	0.01	1.89	18.54	-0.02	-0.39	2.41

Table 5.

Tristimulus Value of Shampoo and Light Fastness Studies with Blue Coupler 2,4-Diaminobenzoxy ethanol on Gray Hair								
3-Aminophenol	PI	Before			After			ΔE
		L*	a*	b*	L*	a*	b*	
Wash 5 hr	1	15.44	0.84	-3.12	17.36	0.39	-3.94	2.14
	2	15.41	0.64	-2.08	17.12	0.79	-3.72	2.37
	3	15.11	0.71	-3.11	18.70	-0.23	-4.89	4.12
Light 72 hour	1	15.44	0.84	-3.12	15.59	-0.17	-1.94	1.56
	2	15.41	0.64	-2.08	15.67	-0.41	-1.19	1.40
	3	15.11	0.71	-3.11	16.51	-0.43	-2.03	2.10

[0070] Results of the experiments are summarized in the following tables 4 and 5. ΔE is defined as


$$\sqrt{(\Delta L^*)^2 + (\Delta a^*)^2 + (\Delta b^*)^2}$$

and indicates the size of the color difference. The smaller the ΔE value, the better the wash and light fastness.

[0071] When coupled with 3-aminophenol, the compound 1 shows better wash fastness property than the compound 3. Total color change (ΔE) of the compound 1 is 0.66, while the compound 3 has ΔE 1.73. Light fastness of 1 is comparable to that of the compound 3 (see Table 4, ΔE : 2.44 vs. 2.41). In the case of blue coupler, 2,4-diaminophenox-yethanol, the compound 1 exhibits better wash and light fastness than the compound 3 (see Table 5).

Claims

1. Use of a compound having the formula I:



wherein R_1 is hydrogen, C_1 - C_6 alkyl and C_1 - C_5 mono or polyhydroxyalkyl, and R_2 is H or OH, or a cosmetically acceptable salt thereof, as a primary dye intermediate in the manufacture of an oxidative hair dye product.

2. Use according to Claim 1, wherein R_1 is hydrogen, C_1 - C_4 alkyl or C_1 - C_3 mono-or polyhydroxyalkyl and R_2 is hydrogen or hydroxyl.

3. Use according to Claim 2, wherein R_1 is hydrogen, methyl or ethyl.

4. Use according to Claim 3 wherein R_1 is hydrogen.

5. Use according to any one of the preceding claims, wherein R_2 is hydrogen.


6. Use according to any one of the preceding Claims wherein said primary dye intermediate is present in an amount of about 0.01% to about 10 %, by weight based on total weight of the oxidative hair dye product.

7. Use according to Claim 6 wherein said primary dye intermediate is present in about 0.1% to about 5%, by weight, based on the total weight of the oxidative hair dye product.

8. Use according to any one of the preceding claims wherein the oxidative hair dye product additionally comprises a coupling compound.

9. Use according to Claim 8 wherein said coupling compound is selected from the group consisting of 3-aminophenol, resorcinol, 2-methylresorcinol, 2-hydroxy-4-aminotoluene, 1-naphthol, 2-(2, 4-diaminophenoxy) ethanol and mixtures thereof.

10. A composition for the oxidative colouring of hair comprising at least one primary dye intermediate; at least one coupling compound, and a cosmetically acceptable vehicle, the primary dye intermediate and the coupling compound forming an oxidative hair dye product in the presence of an oxidising agent, wherein the at least one primary dye intermediate comprises a 1-(4-aminophenyl)-2-pyrrolidinemethanol, or a cosmetically acceptable salt thereof, having the formula I:



wherein R_1 is hydrogen, C_1 - C_6 alkyl and C_1 - C_5 mono or polyhydroxyalkyl, and R_2 is H or OH, and wherein said composition further includes one or more additional dye components selected from the group consisting of: p-phenylenediamine and cosmetically acceptable derivatives thereof; p-aminophenols and cosmetically acceptable derivatives thereof; ortho developers and cosmetically acceptable derivatives thereof; phenols and cosmetically acceptable derivatives thereof; resorcinols and cosmetically acceptable derivatives thereof; m-phenylenediamines and cosmetically acceptable derivatives thereof; m-aminophenols and cosmetically acceptable derivatives thereof; and heterocyclic derivatives.

11. A composition for the oxidative colouring of hair comprising at least one primary dye intermediate; at least one coupling compound, and a cosmetically acceptable vehicle, the primary dye intermediate and the coupling compound forming an oxidative hair dye product in the presence of an oxidising agent, wherein the at least one primary dye intermediate comprises a 1-(4-aminophenyl)-2-pyrrolidinemethanol, or a cosmetically acceptable salt thereof, having the formula I:

5

10

15

20

25

30


35

40

45

50

55



wherein R_1 is hydrogen, C_1 - C_6 alkyl and C_1 - C_5 mono or polyhydroxyalkyl, and R_2 is H or OH, and wherein the composition further includes an oxidising agent.

12. Composition according to Claim 10 or Claim 11 wherein R_1 and R_2 are as defined in any one of Claims 2 to 5.

13. Composition according to any one of Claims 10 to 12 wherein said 1-(4-aminophenyl) -2- pyrrolidinemethanol primary dye intermediate is present in an amount of about 0.01% to about 10 %, by weight based on total weight of the composition.

14. Composition according to Claim 13 wherein said 1-(4-aminophenyl) -2-pyrrolidinemethanol primary intermediate is present in about 0.1% to about 5%, by weight, based on the total weight of the composition.

15. Composition according to Claim 11 further including one or more additional dye components, wherein said additional dye components are selected from the group consisting of: p-phenylenediamine and cosmetically acceptable derivatives thereof; p-aminophenols and cosmetically acceptable derivatives thereof; ortho developers and cosmetically acceptable derivatives thereof; phenols and cosmetically acceptable derivatives thereof; resorcinols and cosmetically acceptable derivatives thereof; m-phenylenediamines and cosmetically acceptable derivatives thereof; m-aminophenols and cosmetically acceptable derivatives thereof; and heterocyclic derivatives.

16. Composition according to any one of Claims 10 to 15, wherein said coupling compound is selected from the group consisting of 3-aminophenol, resorcinol, 2-methylresorcinol, 2-hydroxy-4-aminotoluene, 1-naphthol, 2-(2, 4-diaminophenoxy) ethanol and mixtures thereof.

17. A method for the oxidative colouring of human hair comprising contacting the hair with a hair colouring effective amount of the composition according to any one of Claims 10 to 16 and maintaining contact with the hair until the hair is coloured.

18. A kit comprising in a first container a composition comprising a primary dye intermediate having the formula (I) of Claim 1 and a coupler; and in a second container an oxidising agent.

19. A kit according to Claim 18 wherein the coupler is as defined in Claim 16.

20. A kit according to Claim 18 or Claim 19 wherein said composition is the composition according to Claim 10.

21. A kit according to any one of Claims 18 to 20 wherein said oxidising agent is hydrogen peroxide.

5


Patentansprüche

1. Verwendung einer Verbindung der Formel I:

10

15

20



25

worin R_1 für Wasserstoff, C_1 - C_6 -Alkyl oder C_1 - C_5 - Mono- oder Polyhydroxyalkyl und R_2 für H oder OH steht, oder eines kosmetisch akzeptablen Salzes davon als primäre Farbstoffzwischenverbindung zur Herstellung eines Oxidationshaarfärbemittels.

30

2. Verwendung nach Anspruch 1, wobei R_1 für Wasserstoff, C_1 - C_4 -Alkyl oder C_1 - C_3 -Mono- oder Polyhydroxyalkyl steht und R_2 für Wasserstoff oder Hydroxyl steht.

3. Verwendung nach Anspruch 2, wobei R_1 für Wasserstoff, Methyl oder Ethyl steht.

35

4. Verwendung nach Anspruch 3, wobei R_1 für Wasserstoff steht.

5. Verwendung nach einem der vorhergehenden Ansprüche, wobei R_2 für Wasserstoff steht.

40

6. Verwendung nach einem der vorhergehenden Ansprüche, worin die primäre Farbstoffzwischenverbindung in einer Menge von etwa 0,01 bis etwa 10 Gew.-%, bezogen auf das Gesamtgewicht des Oxidationshaarfärbemittels, vorliegt.

7. Verwendung nach Anspruch 6, worin die primäre Farbstoffzwischenverbindung in einer Menge von etwa 0,1 bis etwa 5 Gew.-%, bezogen auf das Gesamtgewicht des Oxidationshaarfärbemittels, vorliegt.

45

8. Verwendung nach einem der vorhergehenden Ansprüche, wobei das Oxidationshaarfärbemittel außerdem einen Kuppler umfasst.

50

9. Verwendung nach Anspruch 8, wobei der Kuppler ausgewählt ist unter 3-Aminophenol, Resorcinol, 2-Methylresorcinol, 2-Hydroxy-4-aminotoluol, 1-Naphthol, 2-(2, 4-Diaminophenoxy)ethanol und Gemischen davon.


10. Zusammensetzung zur oxidativen Färbung des Haars, umfassend wenigstens eine primäre Farbstoffzwischenverbindung; wenigstens einen Kuppler und einen kosmetisch akzeptablen Träger, wobei die primäre Farbstoffzwischenverbindung und der Kuppler in Gegenwart eines Oxidationsmittels einen Oxidationshaarfärbestoff bilden, worin die wenigstens eine primäre Farbstoffzwischenverbindung ein 1-(4-Aminophenyl)-2-pyrrolidinmethanol der Formel I oder ein kosmetisch akzeptables Salz davon umfasst:

55

5

10

15



20

worin R_1 für Wasserstoff, C_1 - C_6 -Alkyl oder C_1 - C_5 - Mono- oder Polyhydroxyalkyl und R_2 für H oder OH steht, und die Zusammensetzung außerdem eine oder mehrere weitere Farbstoffkomponenten enthält, die ausgewählt sind unter p-Phenylendiamin und kosmetisch akzeptablen Derivativen davon; p-Aminophenolen und kosmetisch akzeptablen Derivativen davon; ortho-Entwicklern und kosmetisch akzeptablen Derivativen davon; Phenolen und kosmetisch akzeptablen Derivativen davon; Resorcinolen und kosmetisch akzeptablen Derivativen davon; m-Phenylendiaminen und kosmetisch akzeptablen Derivativen davon; m-Aminophenolen und kosmetisch akzeptablen Derivativen davon; und heterocyclischen Derivativen.


25

11. Zusammensetzung zur oxidativen Färbung des Haars, umfassend wenigstens eine primäre Farbstoffzwischenverbindung; wenigstens einen Kuppler und einen kosmetisch akzeptablen Träger, wobei die primäre Farbstoffzwischenverbindung und der Kuppler in Gegenwart eines Oxidationsmittels einen Oxidationshaarfärbstoff bilden, worin die wenigstens eine primäre Farbstoffzwischenverbindung ein 1-(4-Aminophenyl)-2-pyrrolidinmethanol der Formel I oder ein kosmetisch akzeptables Salz davon umfasst:

30

35

40



45

worin R_1 für Wasserstoff, C_1 - C_6 -Alkyl oder C_1 - C_5 - Mono- oder Polyhydroxyalkyl und R_2 für H oder OH steht, und die Zusammensetzung außerdem ein Oxidationsmittel enthält.

50

12. Zusammensetzung nach Anspruch 10 oder 11, worin R_1 und R_2 die in einem der Ansprüche 2 bis 5 angegebene Bedeutung haben.

55


13. Zusammensetzung nach einem der Ansprüche 10 bis 12, worin die primäre Farbstoffzwischenverbindung 1-(4-Aminophenyl)-2-pyrrolidinmethanol in einer Menge von etwa 0,01 bis etwa 10 Gew.-%, bezogen auf das Gesamtgewicht der Zusammensetzung, vorliegt.

14. Zusammensetzung nach Anspruch 13, worin die primäre Farbstoffzwischenverbindung 1-(4-Aminophenyl)-2-pyrrolidinmethanol in einer Menge von etwa 0,1 bis etwa 5 Gew.-%, bezogen auf das Gesamtgewicht der Zusammensetzung vorliegt.

15. Zusammensetzung nach Anspruch 11, außerdem enthaltend einen oder mehrere weitere Farbstoffkomponenten, die ausgewählt sind unter p-Phenyldiamin und kosmetisch akzeptablen Derivativen davon; p-Aminophenolen und kosmetisch akzeptablen Derivativen davon; ortho-Entwicklern und kosmetisch akzeptablen Derivativen davon; Phenolen und kosmetisch akzeptablen Derivativen davon; Resorcinolen und kosmetisch akzeptablen Derivativen davon; m-Phenyldiaminen und kosmetisch akzeptablen Derivativen davon; m-Aminophenolen und kosmetisch akzeptablen Derivativen davon; und heterocyclischen Derivativen.
16. Zusammensetzung nach einem der Ansprüche 10 bis 15, worin der Kuppler ausgewählt ist unter 3-Aminophenol, Resorcinol, 2-Methylresorcinol, 2-Hydroxy-4-aminotoluol, 1-Naphthol, 2-(2, 4-Diaminophenoxy)ethanol und Gemischen davon.
17. Verfahren zur oxidativen Färbung des menschlichen Haars, bei dem man das Haar mit einer zur Haarfärbung wirksamen Menge der Zusammensetzung nach einem der Ansprüche 10 bis 16 in Kontakt bringt und im Kontakt mit dem Haar belässt, bis das Haar gefärbt ist.
18. Kit, umfassend in einem ersten Behälter eine Zusammensetzung, die eine primäre Farbstoffzwischenverbindung der Formel I in Anspruch 1 und einen Kuppler umfasst; und in einem zweiten Behälter ein Oxidationsmittel.
19. Kit nach Anspruch 18, worin der Kuppler der Definition in Anspruch 16 folgt.
20. Kit nach Anspruch 18 oder 19, worin es sich bei der Zusammensetzung um eine solche nach Anspruch 10 handelt.
21. Kit nach einem der Ansprüche 18 bis 20, wobei es sich bei dem Oxidationsmittel um Wasserstoffperoxid handelt.

Revendications


1. Utilisation d'un composé ayant la formule I :



- où R_1 est hydrogène, alkyle C_1-C_6 et mono ou polyhydroxyalkyle C_1-C_5 et R_2 est H ou OH, ou son sel cosmétiquement acceptable, en tant qu'intermédiaire de teinture primaire dans la fabrication d'un produit de teinture des cheveux par oxydation.
2. Utilisation selon la revendication 1, où R_1 est hydrogène, alkyle C_1-C_4 ou mono ou polyhydroxyalkyle C_1-C_3 et R_2 est hydrogène ou hydroxyle.
3. Utilisation selon la revendication 2, où R_1 est hydrogène, méthyle ou éthyle.
4. Utilisation selon la revendication 3, où R_1 est hydrogène.
5. Utilisation selon l'une quelconque des revendications précédentes, où R_2 est hydrogène.
6. Utilisation selon l'une quelconque des revendications précédentes où ledit intermédiaire de teinture primaire est

présent en une quantité d'environ 0,01% à environ 10% en poids, en se basant sur le poids total du produit de teinture des cheveux par oxydation.

7. Utilisation selon la revendication 6, où ledit intermédiaire de teinture primaire est présent à raison d'environ 0,1% à environ 5% en poids, en se basant sur le poids total du produit de teinture des cheveux par oxydation.
8. Utilisation selon l'une quelconque des revendications précédentes, où le produit de teinture des cheveux par oxydation comprend additionnellement un composé de couplage.
9. Utilisation selon la revendication 8, où ledit composé de couplage est sélectionné dans le groupe consistant en 3-aminophénol, résorcinol, 2-méthylrésorcinol, 2-hydroxy-4-aminotoluène, 1-naphtol, 2-(2,4-diaminophénoxy)éthanol et leurs mélanges.
10. Composition pour la coloration par oxydation des cheveux comprenant au moins un intermédiaire de teinture primaire ; au moins un composé de couplage et un véhicule cosmétiquement acceptable, l'intermédiaire de teinture primaire et le composé de couplage formant un produit de teinture des cheveux par oxydation en présence d'un agent oxydant où le au moins un intermédiaire de teinture primaire comprend un 1-(4-aminophényl)-2-pyrrolidine-méthanol ou son sel cosmétiquement acceptable ayant la formule I :



- où R_1 est hydrogène, alkyle C_1-C_6 et mono ou polyhydroxyalkyle C_1-C_5 et R_2 est H ou OH, et où ladite composition contient de plus un ou plusieurs composants additionnels de teinture, sélectionnés dans le groupe consistant en p-phénylènediamine et ses dérivés cosmétiquement acceptables ; p-aminophénols et leurs dérivés cosmétiquement acceptables ; développeurs ortho et leurs dérivés cosmétiquement acceptables ; phénols et leurs dérivés cosmétiquement acceptables ; résorcinols et leurs dérivés cosmétiquement acceptables ; m-phénylènediamines et leurs dérivés cosmétiquement acceptables ; m-aminophénols et leurs dérivés cosmétiquement acceptables ; et dérivés hétérocycliques.

11. Composition pour la coloration par oxydation des cheveux comprenant au moins un intermédiaire de teinture primaire ; au moins un composé de couplage et un véhicule cosmétiquement acceptable, l'intermédiaire de teinture primaire et le composé de couplage formant un produit de teinture des cheveux par oxydation en présence d'un agent oxydant, où le au moins un intermédiaire de teinture primaire comprend un 1-(4-aminophényl)-2-pyrrolidineméthanol ou son sel cosmétiquement acceptable ayant la formule I :



où R_1 est hydrogène, alkyle C_1-C_6 et mono ou polyhydroxyalkyle C_1-C_5 et R_2 est H ou OH, et où la composition contient de plus un agent oxydant.

- 20 12. Composition selon la revendication 10 ou la revendication 11, où R_1 et R_2 sont tels que définis dans l'une quelconque des revendications 2 à 5.
- 25 13. Composition selon l'une quelconque des revendications 10 à 12, où ledit 1-(4-aminophényl)-2-pyrrolidinéméthanol comme intermédiaire de teinture primaire est présent en une quantité d'environ 0,01% à environ 10% en poids, en se basant sur le poids total de la composition.
- 30 14. Composition selon la revendication 13, où ledit 1-(4-aminophényl)-2-pyrrolidinéméthanol intermédiaire primaire est présent à raison d'environ 0,1% à environ 5% en poids, en se basant sur le poids total de la composition.
- 35 15. Composition selon la revendication 11 comprenant un ou plusieurs composants additionnels de teinture, où lesdits composants additionnels de teinture sont sélectionnés dans le groupe consistant en p-phénylènediamine et ses dérivés cosmétiquement acceptables ; p-aminophénols et leurs dérivés cosmétiquement acceptables ; développeurs ortho et leurs dérivés cosmétiquement acceptables ; phénols et leurs dérivés cosmétiquement acceptables ; résorcinols et leurs dérivés cosmétiquement acceptables ; m-phénylènediamines et leurs dérivés cosmétiquement acceptables ; m-aminophénols et leurs dérivés cosmétiquement acceptables ; et dérivés hétérocycliques.
- 40 16. Composition selon l'une quelconque des revendications 10 à 15, où ledit composé de couplage est sélectionné dans le groupe consistant en 3-aminophénol, résorcinol, 2-méthylrésorcinol, 2-hydroxy-4-aminotoluène, 1-naphtol, 2-(2,4-diaminophénoxy)éthanol et leurs mélanges.
- 45 17. Méthode pour la coloration par oxydation des cheveux humains comprenant la mise en contact des cheveux avec une quantité efficace de coloration des cheveux de la composition selon l'une quelconque des revendications 1 à 16 et le maintien de contact avec les cheveux jusqu'à ce que les cheveux soient colorés.
- 50 18. Kit comprenant, dans un premier conteneur, une composition comprenant un intermédiaire de teinture primaire ayant la formule (I) de la revendication 1 et un coupleur ; et dans un second conteneur, un agent oxydant.
19. Kit selon la revendication 18 où le coupleur est tel que défini à la revendication 16.
20. Kit selon la revendication 18 ou la revendication 19, où ladite composition est la composition selon la revendication 10.
21. Kit selon l'une quelconque des revendications 18 à 20 où ledit agent oxydant est du peroxyde d'hydrogène.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.